

All questions may be attempted but only marks obtained on the best **five** questions will count.

The use of an electronic calculator is **not** permitted in this examination.

1. (a) Find the radius of convergence of the power series

$$(i) \sum_{n=1}^{\infty} \frac{z^n}{n(n+1)}, \quad (ii) \sum_{n=1}^{\infty} \frac{(-1)^{n+1} z^n}{n^n}.$$

- (b) If z is a complex number, show that $(z + z^{-1})(z - z^{-1}) = z^2 + z^{-2}$. If now $z = \cos \theta + i \sin \theta$, show by expanding $(z + z^{-1})^5(z - z^{-1})^5$ that

$$\sin^5 \theta \cos^5 \theta = \frac{1}{2^9} (\sin 10\theta - 5 \sin 6\theta + 10 \sin 2\theta).$$

- (c) Write down an expression for $\tan z$ in terms of $\exp(iz)$ and use it to find all values of $\tan^{-1}(2i)$.

2. (a) Explain what is meant by the statement “ $f(z)$ is an analytic function in the region D of the complex plane”.
- (b) If $u(x, y)$ and $v(x, y)$ are respectively the real and imaginary parts of an analytic function $f(z)$ with $z = x + iy$, write down the Cauchy-Riemann equations which are satisfied at all points in D and deduce that u and v are harmonic throughout D .
- (c) Show that the analytic function $f(z) = \exp(z)$ has $u(x, y) = \exp(x) \cos y$ as its real part and $v(x, y) = \exp(x) \sin y$ as its imaginary part. Verify the Cauchy-Riemann equations hold everywhere.
- (d) Verify that $h(x, y) = \exp(x)(x \cos y - (y + 1) \sin y)$ is harmonic for all x and y and determine an analytic function $g(z)$ whose imaginary part is $h(x, y)$.

3. (a) Write down Cauchy's integral theorem, integral formula and integral formula for derivatives.
 (b) Find values of A, B, C and D such that

$$f(z) = \frac{z^3 - 1}{z(z^2 - 1)} = A + \frac{B}{z} + \frac{C}{z - 1} + \frac{D}{z + 1}$$

and comment on the result. Find all terms in the Laurent expansions for $f(z)$ in powers of z valid for $|z| < 1$ and for $|z| > 1$.

- (c) Using the results of part (b) or otherwise, find the values of

$$\oint_{C_i} \frac{z^3 - 1}{z(z^2 - 1)} dz,$$

where C_1 is the circle $|z| = 1/2$ and C_2 the circle $|z| = 2$.

4. Use the residue theorem and suitable contours in the complex plane to show

$$(i) \int_0^{2\pi} \frac{d\theta}{5 - 4 \sin \theta} = \frac{2\pi}{3}, \quad (ii) \int_{-\infty}^{\infty} \frac{\cos x \, dx}{x^2 - 2x + 2} = \frac{\pi \cos 1}{e}.$$

5. (a) Euler's equation,

$$\frac{\partial F}{\partial y} - \frac{d}{dx} \left(\frac{\partial F}{\partial y'} \right) = 0,$$

is satisfied by the extremal of the functional

$$\int_a^b F(y, y') dx,$$

in which F has *no explicit dependence on x* , a prime denotes differentiation with respect to x and $y(a)$ and $y(b)$ are prescribed. Show that

$$F - y' \frac{\partial F}{\partial y'}$$

is constant.

- (b) Find the extremal curve of

$$I = \int_0^{\pi/2} (y^2 - (y')^2) \, dx$$

if $y(0) = 0$, $y(\pi/2) = 1$. Show that $I = 0$ on this extremal.

6. The curve $y = y(x)$ is required so that the area

$$I = \int_0^1 y \, dx, \quad y \geq 0,$$

is maximised for a given constant value of the length

$$L = \int_0^1 \sqrt{1 + (dy/dx)^2} \, dx,$$

and subject to the end conditions $y(0) = y(1) = 0$. Show that the extremal satisfies the differential equation

$$\left(\frac{dy}{dx}\right)^2 = \frac{\lambda^2}{(y+c)^2} - 1,$$

for constants λ and c . Show that the solution is a circular arc and derive an equation relating L and the radius of the circular arc.

7. (a) Define the Lagrangian for a conservative system. Write down Lagrange's equations if the system has n degrees of freedom.
- (b) A double pendulum consists of two masses m joined by a light rod AB with the mass at A joined to a fixed point O by a light rod OA . The rods have equal length l and are freely hinged at O and A so that they may move in a vertical plane. If θ_1 and θ_2 are the inclinations of OA and OB to the downward vertical then *you are given* that the potential energy V and kinetic energy T of the pendulum are

$$V = -mgl(2 \cos \theta_1 + \cos \theta_2),$$

$$T = \frac{ml^2}{2}(2\dot{\theta}_1^2 + \dot{\theta}_2^2 + 2\dot{\theta}_1\dot{\theta}_2 \cos(\theta_1 - \theta_2)),$$

where g is the acceleration due to gravity. Show that Lagrange's equations are

$$2\ddot{\theta}_1 + \ddot{\theta}_2 \cos(\theta_1 - \theta_2) + \dot{\theta}_2^2 \sin(\theta_1 - \theta_2) = -2(g/l) \sin \theta_1,$$

$$\ddot{\theta}_1 \cos(\theta_1 - \theta_2) + \ddot{\theta}_2 - \dot{\theta}_1^2 \sin(\theta_1 - \theta_2) = -(g/l) \sin \theta_2.$$

- If the pendulum performs small oscillations about the downward vertical, show that the frequencies, ω , of the normal modes of oscillation satisfy

$$\omega^2 = (2 \pm \sqrt{2})(g/l).$$