

UNIVERSITY COLLEGE LONDON

University of London

EXAMINATION FOR INTERNAL STUDENTS

For The Following Qualifications:–

B.Sc. *M.Sci.*

Mathematics B8: Mathematics For Physics And Astronomy

COURSE CODE : **MATHB008**

UNIT VALUE : **0.50**

DATE : **11-MAY-04**

TIME : **14.30**

TIME ALLOWED : **2 Hours**

All questions may be attempted but only marks obtained on the best **five** solutions will count.

The use of an electronic calculator is **not** permitted in this examination.

1. (a) If $S_n = \sum_{k=0}^{n-1} az^k$, show that $S_n = \frac{a(1-z^n)}{1-z}$ and use this result to show that

$$1 + \cos \theta + \cos 2\theta + \cdots + \cos n\theta = \frac{\cos \left(\frac{n\theta}{2}\right) \sin \left(\frac{(n+1)\theta}{2}\right)}{\sin \left(\frac{\theta}{2}\right)}.$$

- (b) Show that the series

$$\sum_{n=1}^{\infty} \frac{1}{n^2} \left(\frac{z-i}{z+i} \right)^{2n}$$

converges only in the upper-half of the complex plane where $\text{Im}(z) > 0$.

- (c) Write down an expression for $\cos z$ in terms of e^{iz} and use this to find all values of z such that $\cos z = 2$.
2. (a) If a function $f(x + iy) = u(x, y) + iv(x, y)$ is analytic in some region \mathcal{D} of the complex plane, derive the Cauchy-Riemann equations satisfied by u and v in \mathcal{D} . Hence, or otherwise, prove that u and v are harmonic in \mathcal{D} .
- (b) Show that the function $v(x, y) = e^x(x \sin y + y \cos y)$ is harmonic for all x and y . Determine the analytic function $f(z)$, the imaginary part of which is $v(x, y)$ and where $f(0) = i$.

3. (a) Write down Cauchy's Theorem and prove it using Green's Theorem, which states that for some simply connected region R enclosed by a curve C ,

$$\oint_C Pdx + Qdy = \iint_R \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dx dy,$$

for any two continuous functions $P(x, y)$ and $Q(x, y)$.

- (b) Use Cauchy's Integral Formula with the unit circle $|z| = 1$ as the contour chosen in the complex plane to show that, if $f(z)$ is analytic within and on the unit circle, then

$$f(0) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) d\theta.$$

- (c) What is the value of

$$\oint_C \frac{z - i}{z^3 - z^4} dz,$$

if C is

- (i) the circle $|z| = \frac{1}{2}$?
- (ii) the circle $|z + 1| = 3$?

4. Use the residue theorem and a suitable contour in the complex plane to show

$$\int_0^{2\pi} \frac{d\theta}{1 + a \sin \theta} = \frac{2\pi}{\sqrt{1 - a^2}}, \quad \text{for real } a \text{ where } |a| < 1.$$

Using a different contour in the the complex plane, evaluate

$$\int_{-\infty}^{+\infty} \frac{\cos(x)}{(x^2 + b^2)(x^2 + c^2)} dx,$$

where b and c are positive real numbers.

5. (a) If Euler's equation,

$$\frac{\partial F}{\partial y} - \frac{d}{dx} \left(\frac{\partial F}{\partial y'} \right) = 0,$$

is satisfied by the extremal of the functional

$$I = \int_a^b F(y, y') dx,$$

where F has *no explicit dependence on x* , the prime denotes differentiation with respect to x , and $y(a)$ and $y(b)$ are prescribed, show that

$$F - y' \frac{\partial F}{\partial y'}$$

is constant.

- (b) Consider the Brachistochrone problem of finding the curve $y(x)$ down which a particle will slide from the point (x_0, y_0) to the point (x_1, y_1) , where $x_1 > x_0$ and $y_1 < y_0$, in the shortest possible time under gravity (which acts in the negative y -direction). **You are given** that the time T of descent is

$$T = \frac{1}{\sqrt{2g}} \int_{x_0}^{x_1} \frac{\sqrt{1 + (y')^2}}{\sqrt{y_0 - y}} dx.$$

Show that if T is a minimum, the function $y(x)$ satisfies the differential equation

$$(y_0 - y) (1 + (y')^2) = D,$$

for some constant D . Use the substitution $y' = \cot \theta$ to show that

$$\begin{aligned} x &= x_0 + \frac{D}{2} (\sin 2\theta - 2\theta), \\ y &= y_0 - \frac{D}{2} (1 - \cos 2\theta), \end{aligned}$$


is a parametric form of the required curve. How can the constant D be determined?

6. Find the extremal curve for the integral

$$I = \int_0^{\pi/2} (y'^2 + 2xyy') dx,$$

subject to the constraints $y(0) = 0$, $y(\pi/2) = 1$ and $\int_0^{\pi/2} y dx = (\frac{\pi}{2} - 1)$.

7. (a) Define the Lagrangian and Hamiltonian for a conservative system. Write down Lagrange's equations if the system has n degrees of freedom.



- (b) A small bead of mass m slides freely on a smooth uniform circular wire of radius a . The wire is *free* to rotate about its vertical diameter and has moment of inertia of $Ma^2/2$ about the rotation axis. Gravity acts downwards in the vertical direction with acceleration g . The system is described by two generalised coordinates ϕ and θ as shown in the figure, with $0 \leq \theta \leq \pi$.

You are given that the total kinetic energy T and potential energy V of the system are

$$T = \frac{1}{4}Ma^2 \dot{\phi}^2 + \frac{1}{2}m \dot{\phi}^2 a^2 \sin^2 \theta + \frac{1}{2}ma^2 \dot{\theta}^2,$$

$$V = -mga \cos \theta,$$

where a dot indicates a derivative with respect to time.

Use Lagrange's equations to show that

$$\dot{\phi} (M + 2m \sin^2 \theta) \quad \text{and} \quad \frac{1}{4}a^2 \left(M \dot{\phi}^2 + 2m \dot{\phi}^2 \sin^2 \theta + 2m \dot{\theta}^2 \right) - mga \cos \theta$$

are both conserved quantities.

Given the initial state of the system is $\theta = \pi/2$, $\dot{\theta} = 0$ and $\dot{\phi} = \omega$, show that the angular velocity of the wire is

$$\dot{\phi} = \frac{(M + 2m)\omega}{(M + 2m \sin^2 \theta)}.$$

Further, show that the bead does not reach the bottom of the wire ($\theta = 0$) if

$$\omega^2 > \frac{2gM}{a(M + 2m)}.$$