

UNIVERSITY COLLEGE LONDON

University of London

EXAMINATION FOR INTERNAL STUDENTS

For The Following Qualifications:-

B.Sc. *M.Sci.*

Mathematics C325: Real Fluids

COURSE CODE : **MATHC325**

UNIT VALUE : **0.50**

DATE : **07-MAY-04**

TIME : **14.30**

TIME ALLOWED : **2 Hours**

All questions may be attempted but only marks obtained on the best **four** solutions will count. The use of an electronic calculator is **not** permitted in this examination.

You may assume that external (body) forces are absent.

1. Write down the constitutive relation for an isotropic Newtonian fluid with constant density ρ and coefficient of viscosity μ in terms of the stress tensor, σ_{ij} , and the rate of strain tensor, e_{ij} .

Assuming the Cauchy equations of motion,

$$\rho \frac{Du_i}{Dt} = \frac{\partial \sigma_{ij}}{\partial x_j},$$

derive the Navier-Stokes momentum equations for an incompressible Newtonian fluid in the form

$$\rho \left[\frac{\partial \mathbf{U}}{\partial t} + (\mathbf{U} \cdot \nabla) \mathbf{U} \right] = -\nabla p + \mu \nabla^2 \mathbf{U},$$

in the usual notation.

Write down a formula for the rate of energy dissipation per unit volume, ϕ , in terms of the rate of strain tensor, e_{ij} . Calculate the dissipation function ϕ for the steady Couette flow in a planar channel of width d with one wall fixed and the other wall moving with constant speed u_0 .

2. Incompressible fluid of kinematic viscosity ν is contained in a channel between two parallel walls at $y = 0$ and $y = d$ in the (x, y) -plane. The walls of the channel perform periodic oscillations in the x -direction with the speeds $u = u_0 \cos(\omega t)$ and $u = u_1 \cos(\omega t)$, at $y = 0$ and $y = d$, respectively. Here u_0, u_1 are constant and there is no additional pressure gradient along the channel. Verify that a time-periodic, uni-directional, flow is possible with the velocity profile of the form

$$u(y, t) = \operatorname{Re} \left\{ e^{i\omega t} \left[u_1 \frac{\sinh(\alpha y)}{\sinh(\alpha d)} + u_0 \frac{\sinh(\alpha(d-y))}{\sinh(\alpha d)} \right] \right\}$$

with $\alpha = (1 + i) \sqrt{\omega / (2\nu)}$.

Find the limiting form of the velocity profile in the case of large viscosity, $\nu \rightarrow \infty$, and give a qualitative interpretation of your result.

3. Incompressible viscous fluid with density ρ and kinematic viscosity ν flows steadily in an infinitely long planar channel between parallel walls at $y = 0$ and $y = d$ in the (x, y) -plane. A constant pressure gradient, $\partial p / \partial x = -\rho G$, is applied along the channel. The walls are made of a permeable material and the fluid is injected into the channel through the wall at $y = 0$ and extracted from the channel through the second wall at $y = d$ with the same velocity, $v = v_w$, in the y -direction. Show that a two-dimensional flow in the channel is possible with the x -component of the velocity vector of the form

$$u = \frac{G}{v_w} \left[y - d \frac{\exp(v_w y / \nu) - 1}{\exp(v_w d / \nu) - 1} \right].$$

Determine the y -component of the velocity vector in the flow.

In the case of weak injection, $v_w \rightarrow 0$, show that the flow reduces to a Poiseuille flow due to a constant pressure gradient.

Discuss briefly the case of strong suction through the lower wall, $v_w \rightarrow -\infty$, with the focus on the near-wall layer of thickness $y = O(|v_w|^{-1})$.

4. State the assumptions of the lubrication approximation for a two-dimensional flow in a narrow gap between two solid boundaries.

Incompressible viscous fluid fills a narrow gap between a flat plate at $y = 0$ and a flexible wall at $y = h(x, t)$ in the (x, y) -plane. Assuming that the flow is governed by the lubrication approximation, show that

$$h_t = \frac{1}{12\mu} (h^3 p_x)_x$$

where $p = p(x, t)$ is the pressure in the gap.

The pressure along the gap in the range $x > 0$ varies according to

$$p = \frac{p_0}{m+1} x^{m+1}$$

with some constants p_0 and m . Show that the shape of the boundary can take a self-similar form,

$$h(x, t) = t^\alpha H(\xi), \quad \xi = x/t^\beta$$

provided that

$$\alpha H + \frac{1+2\alpha}{m-1} \xi H' = \frac{p_0}{12\mu} (\xi^m H^3)'$$

Derive an implicit solution, $\xi = \xi(H)$ for the case $\alpha = -1, m = 2$ when $\xi > 0$.

5. Show that the streamfunction in a two-dimensional, slow steady flow of incompressible fluid satisfies the equation

$$\nabla^2 (\nabla^2 \psi) = 0.$$

An axisymmetric slow flow is governed by the equation for the streamfunction of the form $D^2 (D^2 \psi) = 0$ where

$$D^2 = \frac{\partial^2}{\partial r^2} + \frac{\sin \theta}{r^2} \frac{\partial}{\partial \theta} \left(\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \right)$$

in spherical polar coordinates. Derive the Stokes solution for the flow past a sphere of radius a placed in a uniform stream with the speed u_∞ .

You may assume the continuity equation

$$\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 u_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (u_\theta \sin \theta) = 0.$$