

UNIVERSITY COLLEGE LONDON

University of London

EXAMINATION FOR INTERNAL STUDENTS

For The Following Qualifications:–

B.Sc. *M.Sci.*

Mathematics C348: Mathematics For General Relativity

COURSE CODE : MATHC348

UNIT VALUE : 0.50

DATE : 19–MAY–06

TIME : 14.30

TIME ALLOWED : 2 Hours

All questions may be attempted but only marks obtained on the best **four** solutions will count.

The use of an electronic calculator is **not** permitted in this examination.

Geodesic Equation ($U^a = dX^a/d\tau$)

$$\frac{dU^a}{d\tau} + \Gamma^a_{bc} U^b U^c = 0; \quad (1)$$

$$\frac{dp_a}{d\tau} = \frac{m}{2} (\partial_a g_{bc}) U^b U^c; \quad (2)$$

$$\Gamma^a_{bc} = \frac{1}{2} g^{ad} (\partial_b g_{cd} + \partial_c g_{bd} - \partial_d g_{bc}). \quad (3)$$

Geodesics parameterized by s

$$\frac{d^2 X^a}{ds^2} + \Gamma^a_{bc} \frac{dX^b}{ds} \frac{dX^c}{ds} = 0. \quad (4)$$

Schwarzschild metric line element

$$d\tau^2 = \left(1 - \frac{r_s}{r}\right) dt^2 - \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2); \quad r_s = 2GM.$$

Faraday tensor

$$F_{ab} = \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & B_z & -B_y \\ E_y & -B_z & 0 & B_x \\ E_z & B_y & -B_x & 0 \end{pmatrix}; \quad F^{ab} = \begin{pmatrix} 0 & E_x & E_y & E_z \\ -E_x & 0 & B_z & -B_y \\ -E_y & -B_z & 0 & B_x \\ -E_z & B_y & -B_x & 0 \end{pmatrix}.$$

Dual Faraday tensor

$$*F^{ab} \equiv \frac{1}{2} \epsilon^{abcd} F_{cd} = \begin{pmatrix} 0 & +B_x & +B_y & +B_z \\ -B_x & 0 & -E_z & E_y \\ -B_y & E_z & 0 & -E_x \\ -B_z & -E_y & E_x & 0 \end{pmatrix}.$$

Maxwell Source Equations

$$\partial_b F^{ab} = j_c^a.$$

Internal Maxwell Equations

$$\partial_a F_{bc} + \partial_b F_{ca} + \partial_c F_{ab} = 0 \quad \text{or} \quad \partial_b *F^{ab} = 0.$$

1. (a) Determine which of the following equations is a valid tensor equation. Describe the errors in the other equations.

(i) $f = G_a{}^b K_a{}^d H^a{}_c L^a{}_d$

(ii) $P_a = A_a{}^b B_b + U_c V^d W_d$

(iii) $X_{ab} = Q^c{}_{bca} + U_b W_a$

(iv) $h = \partial^a V^a - \partial^b \partial_c Z_b{}^c$

- (b) Describe how the Riemann tensor $R^a{}_{bcd}$ transforms from unprimed coordinates X to primed coordinates X' .
- (c) Consider a two-dimensional manifold M with coordinates X^1 and X^2 . Suppose the metric is

$$g_{ab} = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix}$$

Also let V^a be a vector and Q_{ab}, T^{bc} be tensors with values

$$\begin{aligned} V^1 &= 4, & V^2 &= 1; \\ Q_{11} &= 1, & Q_{12} &= 3, & Q_{21} &= 5, & S_{22} &= 7, \\ T^{11} &= 0, & T^{12} &= 2, & T^{21} &= 4, & T^{22} &= 6. \end{aligned}$$

Find the following:

(i) $\bar{\nabla} \cdot \bar{\nabla}$;

(ii) $R^c{}_a = T^{bc} Q_{ba}$.

(iii) $T^a{}_a$.

- (d) Using index notation prove the vector identity

$$\nabla \cdot (\bar{\mathbf{A}} \times \bar{\mathbf{B}}) = \bar{\mathbf{B}} \cdot \nabla \times \bar{\mathbf{A}} - \bar{\mathbf{A}} \cdot \nabla \times \bar{\mathbf{B}}.$$

- (e) Consider a four-dimensional manifold with an anti-symmetric tensor A^{ab} and a symmetric tensor S^{ab} . How many independent components does A^{ab} have? How many independent components does S^{ab} have?
- (f) Suppose a tensor A has components A_E^{ab} in the Earth frame E and components A_S^{cd} in the space frame S . Show that if A_E^{ab} is anti-symmetric then A_S^{cd} will also be antisymmetric.

2. For this question, assume Special Relativity holds (i.e. flat space with $g_{ab} = \eta_{ab}$).
- Write down the Internal Maxwell Equation for $a = 0, b = 2, c = 3$, expressing this equation in terms of the electric and magnetic fields.
 - Show that the Internal Maxwell Equations are trivial (i.e., they give no information) if two of the indices are equal (e.g. if $a = b = 1$).
 - Let ϵ^{abcd} be the totally antisymmetric Levi-Civita tensor. In terms of this tensor, the dual Faraday tensor is defined as $*F^{ab} \equiv 1/2 \epsilon^{abcd} F_{cd}$. Derive a simple expression for $\partial_b *F^{ab}$ from this definition and the Internal Maxwell Equations, showing your work.
 - Express $*F^{ab} F_{ab}$ in terms of \vec{E} and \vec{B} (you may use the expressions for F_{ab} and $*F^{ab}$ given on the first page).
 - In terms of the 4-vector potential ϕ_a , the Faraday Tensor F_{ab} is $F_{ab} = \partial_b \phi_a - \partial_a \phi_b$. The helicity four-vector h^a is defined as

$$h^a = *F^{ab} \phi_b.$$

Show that h^a is conserved ($\partial_a h^a = 0$) if the electric and magnetic fields are everywhere perpendicular.

3. (a) Consider a particle of mass m moving in a geodesic around an object of mass M whose gravitational field is described by the Schwarzschild metric. The particle moves in the plane $\theta = \pi/2$. The Schwarzschild metric in this plane has two symmetries. What are they? Show from the geodesic equation or Noether's theorem that the particle's orbit has two conserved quantities k and h . Express $dt/d\tau$ and $d\phi/d\tau$ in terms of k and h .
- (b) Change variables from r to $u = 1/r$ and parametrize the orbit by ϕ rather than τ . Find the orbit equation for $(du/d\phi)^2$.
- (c) Let $E = mk$ and $L = mh$. From the orbit equation, derive the second order equation

$$\frac{d^2 u}{d\phi^2} = \frac{r_s m^2}{2L^2} - u + \frac{3}{2} r_s u^2.$$

- (d) Consider a photon with $m = 0$. Show that the photon has a circular orbit at

$$u = u_c = \left(\frac{2}{3r_s} \right).$$

Next suppose that a photon is in a nearby orbit, with $u = u_c(1 + \epsilon)$ with initial condition $d\epsilon/d\phi(\phi = 0) = 0$. What is the differential equation for ϵ ? Show that the photon will either spiral in to $r = 0$ or escape to $r = \infty$.

4. Consider a two-dimensional surface embedded in three-dimensional Euclidean space (e.g. the surface of a bowl). Using cylindrical coordinates (r, ϕ, z) , the surface is specified by the function $z = 2\sqrt{r-1}$ for $r > 1$.

(a) Letting $X^1 = r$ and $X^2 = \phi$, show that the metric of this surface is

$$g_{ab} = \begin{pmatrix} r & 0 \\ 0 & r^2 \end{pmatrix}.$$

Also find g^{bc} .

- (b) Calculate the Christoffel symbols Γ^a_{bc} for this metric.
- (c) From the geodesic equation, find the differential equations for a geodesic, i.e. find d^2r/ds^2 and $d^2\phi/ds^2$ for a geodesic parameterized by s .
- (d) Show that the metric has a symmetry, and hence there exists a conserved quantity (call it K) along each geodesic. Express $d\phi/ds$ in terms of K .

5. Consider the surface of the Earth (assumed to be perfectly spherical). In terms of its radius R , co-latitude θ and longitude ϕ , the metric line element is

$$ds^2 = R^2 (d\theta^2 + \sin^2 \theta d\phi^2).$$

- (a) Suppose a ship at some position (θ, ϕ) travels with compass bearing ψ (e.g. $\psi = 0$ if the ship is heading North, and $\psi = \pi/4$ if the ship is heading Northeast). If the ship travels a small distance, with a change $\delta\theta$ in co-latitude and a change $\delta\phi$ in longitude, what is the ratio $\delta\theta/\delta\phi$ in terms of ψ ?
- (b) A Mercator map projection uses coordinates (x, y) , where the coordinate transformations are

$$\begin{aligned} x &= a \phi, \\ y &= a \log \cot \frac{\theta}{2} \end{aligned}$$

with a constant. Find $\delta y/\delta x$ in terms of $\delta\theta/\delta\phi$. Consider a direction on the Mercator map which makes an angle of $\tilde{\psi}$ with respect to the vertical. Show that $\tilde{\psi} = \psi$.

- (c) Find the metric line element ds^2 for the Mercator coordinates x and y (hint: you may use the identity $\sin \theta = 1/\cosh(\log \cot \frac{\theta}{2})$ without proof).
- (d) A *conformally flat* metric in two dimensions has the form

$$g_{ab} = f \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

where f is some function of position. What is the function f for the Mercator map? An angle η between two vectors V^a and W^b can be defined by

$$\cos^2 \eta \equiv \frac{(g_{ab} V^a W^b)^2}{(g_{cd} V^c V^d)(g_{ef} W^e W^f)}.$$

Show that the angle η is independent of the function f .

- (e) A polar map projection uses coordinates $(X^1, X^2) = (\rho, \lambda)$ where

$$\rho = R \sin \theta, \lambda = \phi.$$

What is the metric in these coordinates? Is this metric conformally flat?